

## ***Agrobacterium tumefaciens*-mediated transformation of common bean (*Phaseolus vulgaris*) var. Brunca**

Laura Yesenia Solís-Ramos<sup>1\*</sup>, Julio César Ortiz-Pavón<sup>1</sup>, Antonio Andrade-Torres<sup>2</sup>, Romano Porras-Murillo<sup>1</sup>, Arturo Brenes Angulo<sup>3</sup> & Enrique Castaño de la Serna<sup>4</sup>

1. Biotecnología de Plantas, Escuela de Biología, Universidad de Costa Rica, Sede Rodrigo Facio, San Pedro, San José, Costa Rica; laura.solisramos@ucr.ac.cr; jcesar\_pavon@hotmail.com; rpmurillo@gmail.com
2. Biotecnología y Ecología de Organismos Simbióticos, INBIOTECA (Instituto de Biotecnología y Ecología Aplicada), Universidad Veracruzana, Veracruz, México. CA-UVER-173; aandrade@uv.mx
3. Laboratorio de Biotecnología de Plantas, Centro de Investigaciones Agronómicas (CIA), Universidad de Costa Rica, Sede Rodrigo Facio, San Pedro, San José, Costa Rica; arturo.brenes@ucr.ac.cr
4. Unidad de Bioquímica y Biología Molecular de Plantas, CICY (Centro de Investigación Científica de Yucatán), Mérida, Yucatán, México; enriquec@cicy.mx

\* Correspondence

Received 11-X-2017. Corrected 18-XII-2018. Accepted 17-I-2019.

**Abstract:** Common bean is a crop recalcitrant to *in vitro* regeneration and therefore it lacks an efficient transformation protocol that can be reproduced using *A. tumefaciens*. The main goal of this study was to establish a protocol for *A. tumefaciens* mediated transformation of *Phaseolus vulgaris* var. Brunca by marker genes (*gusA* and *nptII*) together with the gene for trehalose-6-phosphate synthase from *Saccharomyces cerevisiae* (*TPS1*) used in other species to increase tolerance to abiotic stress. The  $\beta$ -glucuronidase activity was detected in 45 % of the LBA4404 ElectroMAX® pCambia1301 infected explants. Transformed explants regenerated new shoots after four to five months period in a kanamycin rich media. Surviving plants were evaluated by PCR and presented an 0.5 % efficiency of transformation. The established protocol for genetic transformation of common bean has two additional advantages with respect to previous reports: (1) it allows for obtaining transformed regenerants and (2) the genetic transformation was stable for the selective gene.

**Key words:** common bean; recalcitrant species; genetic transformation; *A. tumefaciens*; *gusA* activit; *nptII*.

Solís-Ramos, L. Y., Ortiz-Pavón, J. C., Andrade-Torres, A., Porras-Murillo, R., Brenes Angulo, A., & Castaño de la Serna, E. (2019). *Agrobacterium tumefaciens*-mediated transformation of common bean (*Phaseolus vulgaris*) var. Brunca. *Revista de Biología Tropical*, 67(2) Suplemento, S83-S94.

*Phaseolus vulgaris* is cultivated and consumed primarily in Latin America, Africa and Asia (Gepts et al., 2008). Several attempts have been made to transform the species to obtain traits that otherwise cannot be achieved by traditional crosses (Amugune, Anyango, & Mukiyama, 2011). Some genetic manipulations have been carried out with biolistic but at low efficiencies (0.03-0.9 %) (Russell, Wallace,

Bathe, Martinell, & McCabe, 1993; Aragão et al., 1996; Vianna et al., 2004).

Transgenic plants of *P. vulgaris* have been obtained with resistance to bean golden mosaic virus (BGMV) and with increased tolerance to herbicide ammonium glyphosate Imazapyt by the biolistic approach (Aragão, Vianna, Albino, & Rech, 2002; Rech, Vianna, & Aragão, 2008; Aragão, Nogueira, Tinoco, & Faria, 2013).

Transformed plants with the mutant gene associated to the Rep protein have shown a partial resistance to the BGMV and this trait has been then transferred to four commercial cultivars by traditional crosses (Faria et al., 2006). Furthermore a BGMV (EMBRAPA 5.1) was resistant line made by transforming with a genetic silence system against the viral gene rep (AC1) (Bonfim, Faria, Nogueira, Mendes, & Aragão, 2007; Aragão & Faria, 2009; Aragão et al., 2013). However, such transformation has a low efficiency and higher cost compared to the use of *Agrobacterium*-mediated transformation (Amugune et al., 2011).

There have been several efforts to conduct genetic transformation via *A. tumefaciens* (Franklin, Trieu, Cassidy, Dixon, & Nelson, 1993; Zhang, Coyne, & Mitra, 1997), with the efficiency of 2.8- to 28.6 % (Espinosa-Huerta, Quintero-Jiménez, Cabrera-Becerra, & Mora-Avilés, 2013; Collado et al., 2015). Regeneration of plants from transformed explants is difficult in genetic transformation protocols, and for recalcitrant species like common bean is one of the unsolved aspects (Liu, Park, Kanno, & Kameya, 2005; Amugune et al., 2011; Collado et al., 2015). Recently the green nodular compact calli were used as explants to obtain transformed plants, based on the indirect organogenesis regeneration pathway (Collado et al., 2015, 2016).

An efficient system for transformation remains elusive for *Phaseolus vulgaris*. Main goal of this study was to develop a transformation protocol for *Phaseolus vulgaris* var. Brunca via *Agrobacterium tumefaciens* considering several factors like bacterial concentration, incubation of the pre-culture and co-cultivation, antibiotics used and lethal concentrations of selective agents.

## MATERIAL AND METHODS

**Plant materials:** Experiments were performed from 2012 to 2015 using mature common bean seeds var. Brunca (its color black) donated by the Fabio Baudrit Moreno Agricultural Experimental Station [Estación

Experimental Agrícola Fabio Baudrit Moreno (EEAFBM)] of the University of Costa Rica, located in La Garita de Alajuela, Costa Rica.

Seeds were firstly surface-sterilized according to Solís-Ramos et al. (2015). Embryonic axes (EA) were extracted with cutters, tweezers and stereoscopic microscope. The samples were surface-sterilized a second time with a solution of 0.1 % of bleach (commercial sodium hypochlorite) for 10 min and subsequently washed three times with sterile distilled water. Finally, EA were placed in the regeneration medium (RM) in petri dishes. For experiments whose pre-cultivation periods were considered, embryos were extracted at the moment of bacterial immersion by removing cotyledons and radicles.

**Regeneration medium:** Murashige & Skoog (1962) (MS) medium added with 1 mg/L BAP, 30 g/L sucrose, and 6 g/L agar was used (Solís-Ramos et al., 2015). The pH was adjusted to 5.7 prior to autoclave sterilization (120 °C, 15 kg/cm<sup>2</sup>, for 20 min). The cultures were incubated at 26 ± 1 °C with a photoperiod of 12 h light (30 µmol/m<sup>2</sup>s) and 12 h darkness.

**Activation of bacterial strains:** *Agrobacterium* LBA4404-ElectroMAX® cells (transformed with bifunctional BinRD29A or pCAMBIA1301 plasmids) were cultured in 25 mL of liquid Luria-Bertani (LB) medium (10 g/L triptone, 5 g/L NaCl, 5 g/L yeast extract) containing the appropriate antibiotics and allowed to grow for 48 h at 28 °C with 160 rpm agitation according to Solís-Ramos, González-Estrada, Nahuath-Dzib, Zapata-Rodriguez, & Castaño (2009). Subsequently, 200 µl of the bacterial suspension were inoculated in 25 mL of fresh LB medium containing the required antibiotics. The culture was allowed to grow for 24 h at 28 °C and 160 rpm agitation. Then, 25 mL of LB and 100 µM of acetosyringone (4'-hydroxy-3',5'-dimethoxyacetophenone) were added to the 24-h culture which was incubated for 4 hr at 28 °C and 160 rpm agitation. The bacterial suspension was centrifuged at 5 000 rpm during 5 min at room temperature

and resuspended in 20 mL of MS liquid media containing 1 mg/L BAP and 100  $\mu$ M of acetosyringone until it reached 0.5 or 0.8 OD<sub>600nm</sub> depending on the treatment. The inoculation was performed by immersing the explants in the bacterial suspension for 15 min under constant agitation (900 rpm), and bacterial suspension excess was eliminated with sterile towel paper. The co-cultivation was performed in Petri dishes with regeneration medium under complete darkness at 25 $\pm$ 1 °C and co-cultivation time depended on the treatments of each experiment.

**Bacteria elimination after co-cultivation:** The *Agrobacterium* elimination was performed after the co-cultivation in pre-cultivated or non-pre-cultivated explants. Explants were shaken at 90 rpm for 15 min in the bacterial suspension and then co-cultivated for four days. After co-cultivation, explants were washed three times with sterile distilled water for 5-10 min (manual agitation) with subsequent vacuum infiltration at 1.25 plg Hg during 30-40 min with solutions containing timentin, cefotaxime or a mixture of both. The following treatments were evaluated: (T1) non-infected EA on antibiotic-free RM, (T2) co-cultivated EA on antibiotic-free RM, (T3) cocultivated EA on RM containing 500 mg/L timentin and 500 mg/L cefotaxime, (T4) co-cultivated EA on RM containing 500 mg/L timentin, and (T5) co-cultivated EA on RM containing 750 mg/L timentin.

EAs water excess was adsorbed with sterile paper towels before they were cultured in the RM supplemented with the selected treatment. Then, they were cultured under the conditions described previously. Twelve explants per treatment were used and the bacterial overgrowth (expressed as percentage) after seven days of cultivation was calculated. The best treatment to eliminate bacteria was selected and its effect on the *in vitro* regeneration was assessed.

**Assessment of kanamycin (Kn) as selective agent:** The explants with a pre-cultivation time of seven days were cultured in RM

supplemented with 0, 25, 50, 100, 200 and 300 mg/L of kanamycin. Each treatment was replicated three times with seven explants per treatment. The explants were cultured at 25 °C under a 16 hr light/8 hr dark photoperiod. Sub-culture was performed every three weeks to the same medium. After three and six weeks of culture we recorded the survival rate, as well as the number of shoots and/or roots observed per explant.

#### **Transient transformation experiments:**

The *A. tumefaciens* LBA4404-ElectroMAX® strain carrying the binary vector pCAM-BIA1301 (11837 bp) was used. This vector contains the  $\beta$ -glucuronidase (*gusA*) reporter gene of *Escherichia coli* with an intron. The *gusA* gene under the 35S promoter of the cauliflower mosaic virus (CaMV) and the poly(A) ending sequence contains the gene *hpt* (hygromycin phosphotransferase, a selective marker gene) controlled by the 35S promoter and the poly (A) 35S CaMV terminator.

Non-precultured EAs were incubated at room temperature in the bacterial suspension under 0.5 OD<sub>600nm</sub>. Later, bacterial excess was adsorbed with sterile filter paper and EAs were placed in Petri dishes containing RM. The effect of the co-culture period was assessed under three, four, and five days of incubation at 25 °C in the dark (Table 1). A second experiment assessed a pre-culture time of two days and a co-culture time of four days (Table 1).

**Histochemical staining:** After bacterial elimination, the *gusA* activity on EAs was detected by histochemical staining (Jefferson, Kavanagh, & Bevan, 1987). Explants were incubated overnight at 37 °C in 100 mM of phosphate sodium (NaH<sub>2</sub>PO<sub>4</sub>, pH 7.0) buffer containing 10 mM EDTA, 0.5 mM potassium ferrocyanide K<sub>3</sub>Fe (CN)<sub>6</sub>, 0.5 mM potassium ferrocyanide K<sub>3</sub>Fe (CN)<sub>6</sub>·3H<sub>2</sub>O, and 0.2 % Triton X-100. X-Gluc (5-bromo-4-chloro-3-indolyl- $\beta$ -D-glucuronide) dissolved in dimethyl sulfoxide (DMSO) under a final concentration of 0.7 mg/L was added to the solution. After incubation, the staining solution

TABLE 1  
Conditions of the genetic transformation of bean with LBA4404-ElectroMAX® of *Agrobacterium*

Experiment	Pre-cultivation period (Days)	Number of explants	Plasmid	Infection	Co-cultivation period (Days)
1	0	11	pCAMBIA1301	Immersion and agitation for 15 min.	3
		11			4
		11			5
2	2	197	pCAMBIA1301	Vacuum infiltration, 1 h, 1.25 plg Hg	4
3	12	30	pCAMBIA1301	Immersion and agitation for 15 min	3
4	2	405	pBinRD29A-bifuncional	Vacuum infiltration, 1 h, 1.25 plg Hg	4
5	8	60	pBinRD29A-bifuncional	Vacuum infiltration, 1 h, 1.25 plg Hg	4
6	10	120	pBinRD29A-bifuncional	Vacuum infiltration, 1 h, 1.25 plg Hg	4
7	12	30	pBinRD29A-bifuncional	Immersion and agitation for 15 min	3
8	17	45	pBinRD29A-bifuncional	Vacuum infiltration, 1 h, 1.25 plg Hg	4

was removed and EAs were put in ethanol (70 %) for 24 h to eliminate chlorophylls. The presence of blue points was identified by means of a stereomicroscope (LEICA E24HD) and was registered and interpreted as transient expression of *gusA*. Photographs were taken with the digital camera of the stereomicroscope (LEICA E24HD). The transient expression of *gusA* was measured by counting the number of explants with at least one blue foci. The frequency of transient expression of GUS was calculated as the number of explants with at least one blue foci over the total number of explants (expressed as percentage).

**Stable transformation:** To assess the stability of the transformation binary and bifunctional vector pBindrd29A was used. This vector contains the selective marker *nptII*, which provides resistance to kanamycin and a fusion of genes containing regions of trehalose-6-phosphate synthase (*tps1*) and trehalose-6-phosphatase synthase (*tps2*) from *Saccharomyces cerevisiae* (Miranda et al., 2007) under the stress inducible promoter rd29A of *Arabidopsis thaliana* (Yamaguchi-Shinozaki & Shinozaki, 1994). This vector was added to the *Agrobacterium tumefaciens* LBA4404-ElectroMAX® strain. The vector and the fusion of the bifunctional genes (rd29A::tps-tps) had a size of 16.02 and 2.9 kb, respectively (Santamaría et al., 2009). Bacterial preparation was performed

as indicated previously. Transformation of common bean was carried out during two, eight, ten, 12, and 17 days of precultivation, and three and four days co-culture (Table 1).

After the co-cultivation period, bacteria were eliminated with the selected treatment of previous experiments. Explants were then cultured to RM added with 500 mg/L of timentin and 50 mg/L of kanamycin under the conditions previously described. Sub-cultures were made every three - four weeks for more than 120 days. The number of shoots surviving the selection process since the first sub-culture, was recorded. The transformation frequency of the selective gene was calculated as the number of kanamycin-resistant explants over the total number of explants.

**PCR amplification of *nptII* and *TPSI* genes:** Genomic DNA was isolated from leaf tissues from four-five month old regenerated bean plants according to the extraction kit protocol MACHERY-NAGEL.

The extraction of DNA from pBindrd29A was performed according to the manual for plasmid purification (INVITROGEN). The primers used for PCR amplification correspond to the *TPSI* sequence forward (5' GTG-GCAGAGGAGCTTGTTGAG 3'), and reverse (5' GGTACTCACATACAGAC 3'), which amplify a fragment of approximately 1600 bp. The final composition of the reaction mixture

was: 2.5 µl buffer 10 X, 1.0 µL Nucleotide Mix 10mM, 2 µl MgCl<sub>2</sub> 50 mM, 1 µL Forward primer (stock 10 picomol/µl) and 1 µL reverse primer (stock 10 picomol/µL), 1 µl plasmid DNA and 0.2 µl taq. The reaction conditions were: 94 °C for 5 min (1x), 30 amplification cycles (94 °C for 1 min, 50 °C for 1 min, 72 °C for 2 min), and a final extension step at 72 °C for 10 min.

The primers for the *nptII* gene were: forward 5'-GAGGCTATTCGGCTATGACTG and reverse 5'-TCGACAAGACCGGCTTC-CATC (Aragão et al., 1996), which amplify a 410 bp fragment. The final composition of the reaction mixture was: 10 µl Master Mix 10mM, 0.25 µl forward primer (stock 10 µM/µl) and 0.25 µl reverse primer (stock 10 picomol/µl), 1 µl DNA (1:10). The reaction conditions were 94 °C for 5 min (1x), 35 amplification cycles (35 cycles of 94 °C for 30 s, 58 °C for 30 s and 72 °C for 1 min) and a final step at 72 °C for 5 min and 4 °C.

PCR products were separated by electrophoresis on 1 % (w/v) agarose gels under 100 V for 50 min and stained with Loading Dye 6X (added with 0.6 % of Gel Red) (8 µl of PCR product was mixed with 2 µl of Loading Dye 6X 2µl molecular marker weight, 2 µl Gel Red. Gene Ruler 1 Kb Plus DNA Ladder, ready to use).

**Presence of *A. tumefaciens* in transformed plants:** The amplification by PCR of VirE2 gene fragment of *A. tumefaciens* was examined using the primers: forward

(5'-TGCCCACCAAGGCGGAATT-3') and reverse (5'-CTTTGCCGACCCATCGA-3') which amplifies a fragment of 895 bp. The volume of the reaction mixture was of 25 ml containing 2.5 µl buffer 10X, 50 mM MgCl<sub>2</sub>, 10 mM dNTPS, 0.1 µl Taq polymerase, 1 µl sense primer, 1 µl antisense primer and 1 µl plant DNA. The reaction conditions were 94 °C for 30 seconds, 30 cycles of temperature of denaturing 94°C for 5 min, primer annealing 55°C for 5 min, elongation 72°C for 2 min and 72 °C for 10 min and 4 °C (10 min). The amplified fragments were separated in 1 % (w/v) agarose gel to 100 V by 40 min.

All the experiments were undertaken under completely randomized designs and the statistical analysis was performed using one-way ANOVA. Means (shoots per explants, height of shoots, roots per regenerants and length of roots) were separated by the Duncan's test at *P* < 0.05. The program SPSS Statistics version 23 was used (IBM Corp., 2015).

## RESULTS

***A. tumefaciens* elimination after the co-culture:** Pre-cultured and co-cultured explants with LBA4404-ElectroMAX® *Agrobacterium* showed a 100 % bacterial suppression regardless the concentrations of cefotaxime and timentin. However, the same antibiotics treatments did not completely eliminate bacteria in non-pre-cultured explants (Table 2).

For the elimination of *A. tumefaciens* the most efficient and compatible with further

TABLE 2  
Recidivism of LBA4404-ElectroMAX® *A. tumefaciens* in explants of *P. vulgaris* L.  
with or without pre-cultural times and co-cultivated during four days

Treatments	% of Bacterial Recidivism	
	Without pre-culture	With pre-culture
Control (with neither co-culture nor antibiotics)	0	0
Control (with co-culture and RM without antibiotics)	92	100
RM+500 mg/L timentin + 500 mg/L cefotaxime	67	0
RM+500 mg/L timentin	92	0
RM+ 750 mg/L timentin	92	0

Observations were taken after six days of starting the treatment.




development was the use of 500 mg/L timentin and also in the RM did not affect the in vitro response of the common bean (data not shown).

**Assessment of the kanamycin as selective agent:** Among the assessed Kn concentrations, 25 mg/L was the only concentration that did not caused complete mortality of the explants after six weeks in culture, while concentrations of around (or higher than) 50 mg/L reached a 100 % mortality. Hence, the following experiments used a minimum lethal dose of 50 mg/L Kn to select the bean cells which were possibly transformed by the *nptII* gene.

**Transient transformation experiments:** Non-pre-cultured EAs inoculated with 0.5 OD<sub>600nm</sub> and co-cultured for three and five days showed  $\beta$ -glucuronidase activity in 45 % and 1.4 % of total explants, respectively

(Fig. 1A). In the case of EAs co-cultured for four days, no  $\beta$ -glucuronidase activity was detected. Only 0.5 % of the explants pre-cultured for two days and co-cultured for four days showed *gusA* activity (Fig. 1 C). Furthermore, non-co-cultured explants showed an absence of endogenous GUS activity (Fig. 1B and Fig. 1D).

**Stable genetic transformation:** Regenerants from 50 mg/L kanamycin treatment, were obtained from explants co-cultured with pBindrd29A (Table 1). These regenerants were selected 120 days after the first sub-culture (Table 3). A higher number of regenerants in the presence of the selective medium was observed in those cases under a long-term pre-culture (equal or higher than eight days). Regenerants from explants pre-cultured for 12 days and co-cultured for three days showed a



**Fig. 1.** *GusA* activity in *P. vulgaris* L. explants co-cultivated with *A. tumefaciens* LBA4404 pCambia1301 with 0.5 OD<sub>600nm</sub>. A and B: non pre-cultured. C and D: pre-cultured for two days. A. co-cultivated EA for five days and B. non-infected EA (negative control), C. co-cultivated EA for four days and D. non-infected EA. Scale bars A and B= 2 mm, C and D= 5 mm.

higher survival rate (Table 3). Explants with a short-term pre-culture (two days) showed a low transformation frequency for the selective agent (Table 3). Fig. 2A indicates the presence of *A. tumefaciens* cells. In contrast, kanamycin-based selected plants did not show this amplification and the presence of the *nptII* gene was detected (Fig. 2B).

**PCR assessment of possible transformation:** Plants pre-cultured and co-cultured for two and four days respectively, showed an amplification fragment of a 410 bp which corresponded to the expected size for the selective gene (Fig. 2A). Kanamycin-based selected plants did not show this amplification. Regenerants with kanamycin resistance were assessed by PCR to determine the presence of the *nptII* gene (Fig. 2 B). These results indicate a transformation efficiency of 0.5 % for the *nptII* gene in plants of common bean.

### DISCUSSION

The main limitation/constrain for obtaining transformed bean plants is the lack of an adequate genetic transformation system due to recalcitrance to the *in vitro* regeneration and the low *Agrobacterium*-mediated transformation rates.


Bacteria elimination after co-culture was also very difficult as the only way to genetically transform beans by the time this work started was by biolistics (Faria et al., 2006). Results indicated that 500 mg/L timentin eliminated the LBA4404 ElectroMax strain of *Agrobacterium tumefaciens* after the co-culture, which is in accordance to recent reports (Mukeshimana, Ma, Walworth, Song, & Kelly, 2013).

Bean EAs co-cultured for three days under 0.5 OD600nm and incubated at 25 °C showed β-glucuronidase activity. This suggests a transient transformation of 45 %, which is similar to that recently reported by Collado et al. (2016), (46 %) using the C58C1RifR strain and a higher co-culture time (five days). This transformation efficiency increased (80 %) with the EHA101 strain under the same conditions

TABLE 3  
Morphologic traits and frequency of regenerants of beans co-cultured with LBA4404-ElectroMAX® pBIndrd29A and selected with 50 mg/L kanamycin after more than 120 days

Experiment	Pre-culture period (days)	Co-culture period (days)	Selection period (days)	Selected nptII regenerants ( %)	Shoots per explants (Average)	height of shoots cm (Average)	Explants with roots (%)	Roots per regenerant (Average)	Length of roots cm (Average)
4	2	4	120	0.5	2.50 ± 2.12	4.25 ± 3.88	0	0 ± 0	0 ± 0
5	8	4	133	16	0.10 ± 0.31	0.10 ± 0.31	10	0.20 ± 0.63	0.30 ± 0.94
6	10	4	133	19	1.04 ± 1.19	0.79 ± 1.09	21.73	0.46 ± 1.27	1.13 ± 3.45
7	12	3	148	36	0.91 ± 0.70	1.00 ± 1.20	27.27	0.69 ± 1.54	1.19 ± 3.32
8	17	4	133	8	0 ± 0	0 ± 0	0	0 ± 0	0 ± 0

No significant differences between means (P < 0.05).



**Fig. 2.** Plants of common bean transformed with the *nptII* gene (120 days of culture). **A.** Amplification of a 410 bp fragment corresponding to the expected size of the selective gene. **B.** Absence fragment *virE* in transformed tissues, line 1: DNA ladder (1 Kb), line 2: *Agrobacterium*, Lines 3-4: regenerated and selected plants (third experiment) and line 5: load control. **C.** Regenerated and kanamycin-resistant plants. **D.** Non-transformed plants. Scale C and D bars = 1 cm.

(Collado et al., 2016). These results evidence the lack of endogenous *gusA* activity in the assessed bean explants. However, this is not the case for other species (Solís-Ramos, González-Estrada, Andrade-Torres, Godoy-Hernández, & Castaño de la Serna, 2010). The *gusA* reporter gene is useful for experiments related to the transformation of certain species and is suitable for histochemical staining and localization of transformed cells.

In this work, the regeneration or “shoots proliferation” from EAs was consistent with previous reports that concluded this type of explant is regenerable but not optimum for the *Agrobacterium*-mediated genetic transformation and subsequent shoots production (Delgado-Sánchez et al., 2006; Arellano, Fuentes, Castillo-España, & Hernández, 2009;

Quintero-Jiménez, Espinosa-Huerta, Acosta-Gallegos, Guzmán-Maldonado, & Mora-Avilés, 2010; Mukeshimana et al., 2013; Solís-Ramos et al., 2015).

Optimum bacteria concentration is a relevant parameter for a transformation protocol (Collado et al., 2015). An OD<sub>600nm</sub> lower than 0.5 will not allow for genetic transformation and higher concentrations kill the explants. According to previous results (data not shown), the bacteria (LBA4404-ElectroMAX®) were used at an OD<sub>600nm</sub> of 0.5. Previous reports used similar conditions (Mukeshimana et al., 2013; Collado et al., 2015, 2016).

The OD<sub>600nm</sub> was combined with short-term co-cultures (three - four days). Three-days co-cultures favored the insertion of the marker gene (*GUS* or *nptII*), which agrees with the



results reported by (Amugune et al., 2011). On the other hand, previous studies indicate long-term co-cultures of five days (Collado et al., 2016), six days (Collado et al., 2015), and eight days (Espinosa-Huerta et al., 2013; Mukeshimana et al., 2013) increase the temporary expression of the transgene (Zambre et al., 2003; Mukeshimana et al., 2013) especially in embryonic axes (Mukeshimana et al., 2013). The presence of BAP in the co-culture and regeneration media favored transformation and survival of the explants. Other studies report similar results (Zhang et al., 1997; Amugune et al., 2011).

Unlike other species, bean did not show natural resistance to kanamycin (Solís-Ramos et al., 2009) as 50 mg/L allowed for selecting the bean regenerants for a four-month time period. In this study, the selection was applied after the first sub-culture (four weeks) to allow transformed cells to proliferate and develop a group of cells capable to survive in middle of a mass of death cells derived from a lethal dose of antibiotic (Zambre et al., 2005). According to Amugune et al. (2011), it should be possible to recover *Agrobacterium*-transformed plants after bean explants had been exposed to the selection medium during three to four weeks (one month). Among all the shoots selected during more than 120 days, only four of them showed an amplification corresponding to the expected size of the *nptII* gene. These results evidence that the type of the explant (embryonic axes) used in the experiment is not suitable for the regeneration of transformed plants (Mukeshimana et al., 2013), as the obtained shoots are from multicellular origin, which may inhibit the strict selection of transgenic shoots, and allows a high number of “leaks” (non-transgenic plants that survive the selection) and chimeras (Angenon & Thu, 2011). Therefore, an important contribution to enhance bean is the establishment of a regeneration system via somatic embryogenesis, or a regeneration system derived from the meristem-free tissue that minimize the production of chimeric transformants (Mukeshimana et al., 2013).

In this study, the transformation frequency obtained for the reporter and selective gene was 0.5 %, which is slightly higher to that reported for the species using biolistic 0.3 % (Aragão et al., 1996; Faria et al., 2006; Rech et al., 2008). Recently, higher transformation efficiencies (2.8 and 0.75 %) have been reported using hypervirulent strains of *A. tumefaciens* (EHA105 or EHA101). This affects the transfer capacity of T-DNA as chromosomal and virulence genes may differ from one strain to another (Collado et al., 2015, 2016). In the case of the *TPS1* gene, the fragment was amplified in transformed explants and the controls explants, and it was not possible make a conclusion on this gene.

So far, few studies report the *Agrobacterium*-assisted genetic transformation for bean and no plants have been obtained, due to the lack of a regeneration protocol (Angenon & Thu, 2011; Mukeshimana et al., 2013). However, Collado et al. (2015, 2016) could recently transform beans with genetic markers (*bar*, *nptII* y *uidA*) producing chimeric regenerants through direct and indirect organogenesis from epicotyl. However, the established protocol did not rendered stable genetic transformation (Collado et al., 2016).

In this study, explants of common bean co-cultivated with LBA4404-ElectroMAX® from *A. tumefaciens*, were obtained. These showed β-glucuronidase activity indicating a temporary genetic transformation (*gusA* gene) of 45 % efficiency. Bean regenerants with kanamycin were selected during three to five weeks for stable transformation. However, this selection was not strict because the regenerants from multicellular origin generated chimeric transformants, leading to leaks of non-transgenic materials. Stable transformation for the *nptIII* gene was obtained at 0.5 % efficiency, which agrees with the results related to the temporary transformation for *gusA* using the same protocol. The established protocol for genetic transformation of common bean has two additional advantages with respect to previous reports: (1) it allows for obtaining transformed regenerants

and (2) the genetic transformation was stable for the selective gene.

**Ethical statement:** authors declare that they all agree with this publication and made significant contributions; that there is no conflict of interest of any kind; and that we followed all pertinent ethical and legal procedures and requirements. All financial sources are fully and clearly stated in the acknowledgements section. A signed document has been filed in the journal archives.

## ACKNOWLEDGEMENTS

Thanks to the Fabio Bautrit Moreno Agricultural Experimental Station (EEAFBM-UCR) and Juan Carlos Hernández (UCR-PITTA bean) for providing the variety of common bean used in this research. Also, thanks to Norman Rojas (Escuela de Microbiología, UCR) for providing the *Escherichia coli* (HB101) strain, Wilma González Kantún (Bioquímica y Biología Molecular de Plantas, CICY, Mérida, Yucatán) and Bruce Williamson Benavides for the technical support, Carolina Céspedes Garro and Andrea Ramírez Telles (Escuela de Biología, UCR) for repeating the molecular assessments of the potential regenerated and selected transformants, and to the DAAD for additional funding. Finally, thanks to the Vice-Rector of Research (Universidad de Costa Rica), for research funding (project No. 111-B2-052). To the CICY (Centro de Investigación Científica de Yucatán) for the vectors donated for this research.

## RESUMEN

**Transformación de frijol común (*Phaseolus vulgaris*) var. Brunca mediada por *Agrobacterium tumefaciens*.** El frijol común en un cultivo recalcitrante a la regeneración *in vitro* y se carece de un protocolo eficiente y reproducible de transformación genética usando *A. tumefaciens*. Desarrollamos un protocolo de transformación genética mediada por *A. tumefaciens* de frijol común variedad Brunca utilizando genes marcadores (*gusA* y *nptII*) junto con el gen de la trehalosa-6-fosfato sintasa de levadura (*TPS1*) utilizado para incrementar tolerancia a estrés

abiótico. La actividad de la  $\beta$ -glucuronidasa fue detectada en 45 % de los explantes infectados con la cepa LBA4404 de *A. tumefaciens* transformada con pCambia1301. Después de 4 o 5 meses se regeneraron tallos en un medio adicionado con kanamicina. Los explantes supervivientes se evaluaron mediante PCR y presentaron una eficiencia de transformación de 0.5 %. El protocolo de transformación genética de frijol común establecido tiene dos ventajas adicionales con respecto a los reportes previos: (1) permite la obtención de regenerares transformados y (2) la transformación genética fue estable para el gen selectivo.

**Palabras clave:** frijol común; especies recalcitrantes; transformación genética; *A. tumefaciens*; actividad *gusA*; *nptII*.

## REFERENCES

- Amugune, N. O., Anyango, B., & Mukiyama, T. K. (2011). *Agrobacterium*-mediated transformation of common bean. *African Crop Science Journal*, 19(3), 137-147.
- Angenon, G., & Thu, T.T. (2011). Genetic transformation. In A. Pratap & J. Kumar (Eds.), *Biology and Breeding of Food Legumes* (pp. 178-192). USA: CABI.
- Aragão, F. J. L., Barros, L. M. G., Brasileiro, A. C. M., Ribeiro, S. G., Smith, F. D., Sanford, J. C., Rech, E. L. (1996). Inheritance of foreign genes in transgenic bean (*Phaseolus vulgaris* L.) co-transformed via particle bombardment. *Theoretical and Applied Genetics*, 93(1-2), 142-150. DOI: 10.1007/BF00225739
- Aragão, F., & Faria, J. (2009). First transgenic geminivirus-resistant plant in the field. *Nature Biotechnology*, 27, 1086-1088. DOI:10.1038/nbt1209-1086
- Aragão, F. J. L., Nogueira, E. O. P. L., Tinoco, M. L. P., & Faria, J. C. (2013). Molecular characterization of the first commercial transgenic common bean immune to the Bean golden mosaic virus. *Journal of Biotechnology*, 166(1), 42-50. DOI: 10.1016/j.jbiotec.2013.04.009
- Aragão, F. J. L., Vianna, G. R., Albino, M. M. C., & Rech, E. L. (2002). Transgenic Dry Bean Tolerant to the Herbicide Glufosinate Ammonium. *Crop Science*, 42(4), 1298. DOI: 10.2135/cropsci2002.1298
- Arellano, J., Fuentes, S. I., Castillo-España, P., & Hernández, G. (2009). Regeneration of different cultivars of common bean (*Phaseolus vulgaris* L.) via indirect organogenesis. *Plant Cell, Tissue and Organ Culture*, 96(1), 11-18. DOI: 10.1007/s11240-008-9454-1
- Bonfim, K., Faria, J. C., Nogueira, E. O. P. L., Mendes, É. A., & Aragão, F. J. L. (2007). RNAi-Mediated Resistance to Bean golden mosaic virus in Genetically Engineered Common Bean (*Phaseolus vulgaris*). *Molecular Plant-Microbe Interactions*, 20(6), 717-726. DOI: 10.1094/MPMI-20-6-0717

- Collado, R., Bermúdez-Caraballosa, I., García, L. R., Veitia, N., Torres, D., Romero, C., & Angenon, G. (2016). Epicotyl sections as targets for plant regeneration and transient transformation of common bean using *Agrobacterium tumefaciens*. *In Vitro Cellular & Developmental Biology - Plant*, 52(5), 500-511. DOI: 10.1007/s11627-016-9769-2
- Collado, R., Bermúdez-Caraballosa, I., García, L. R., Veitia, N., Torres, D., Romero, C., Angenon, G. (2015). *Agrobacterium*-mediated transformation of *Phaseolus vulgaris* L. using indirect organogenesis. *Scientia Horticulturae*, 195 (Supplement C), 89-100. DOI: 10.1016/j.scienta.2015.06.046
- Delgado-Sánchez, P., Saucedo-Ruiz, M., Guzmán-Maldonado, S. H., Villordo-Pineda, E., González-Chavira, M., Fraire-Velázquez, S., ... Mora-Avilés, A. (2006). An organogenic plant regeneration system for common bean (*Phaseolus vulgaris* L.). *Plant Science*, 170(4), 822-827. DOI: 10.1016/j.plantsci.2005.11.015
- Espinosa-Huerta, E., Quintero-Jiménez, A., Cabrera-Becerra, K. V., & Mora-Avilés, M. A. (2013). Stable and efficient *Agrobacterium tumefaciens* mediated transformation of *Phaseolus vulgaris*. *Agrociencia*, 47(4). Retrieved from <http://www.redalyc.org/resumen.oa?id=30226975002>
- Faria, J. C., Albino, M. M. C., Dias, B. B. A., Cançado, L. J., da Cunha, N. B., de M. Silva, L., ... Aragão, F. J. L. (2006). Partial resistance to Bean golden mosaic virus in a transgenic common bean (*Phaseolus vulgaris* L.) line expressing a mutated rep gene. *Plant Science*, 171(5), 565-571. DOI: 10.1016/j.plantsci.2006.06.010
- Franklin, C. I., Trieu, T. N., Cassidy, B. G., Dixon, R. A., & Nelson, R. S. (1993). Genetic transformation of green bean callus by *Agrobacterium* mediated DNA transfer. *Plant Cell Reports*, 12(2), 74-79. DOI: 10.1007/BF00241938
- Gepts, P., Aragão, F. J. L., De Barros, E., Blair, M. W., Brondani, R., Broughton, W., Yu, K. (2008). Genomics of Phaseolus Beans, a Major Source of Dietary Protein and Micronutrients in the Tropics. In P. H. Moore, & R. Ming (Eds.), *Genomics of Tropical Crop Plants* (pp. 113-143). New York: Springer. DOI: 10.1007/978-0-387-71219-2\_5
- IBM Corp. (2015). *IBM SPSS Statistics for Windows* (Version 23.0). Armonk, NY: IBM Corp.
- Jefferson, R. A., Kavanagh, T. A., & Bevan, M. W. (1987). GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *The EMBO Journal*, 6(13), 3901-3907.
- Liu, Z., Park, B.-J., Kanno, A., & Kameya, T. (2005). The Novel Use of a Combination of Sonication and Vacuum Infiltration in *Agrobacterium*-mediated Transformation of Kidney Bean (*Phaseolus vulgaris* L.) with lea Gene. *Molecular Breeding*, 16(3), 189. DOI: 10.1007/s11032-005-6616-2
- Miranda, J. A., Avonce, N., Suárez, R., Thevelein, J. M., Dijck, P. V., & Iturriaga, G. (2007). A bifunctional TPS-TPP enzyme from yeast confers tolerance to multiple and extreme abiotic-stress conditions in transgenic *Arabidopsis*. *Planta*, 226(6), 1411-1421. DOI: 10.1007/s00425-007-0579-y
- Mukeshimana, G., Ma, Y., Walworth, A. E., Song, G., & Kelly, J. D. (2013). Factors influencing regeneration and *Agrobacterium tumefaciens*-mediated transformation of common bean (*Phaseolus vulgaris* L.). *Plant Biotechnology Reports*, 7(1), 59-70. DOI: 10.1007/s11816-012-0237-0
- Murashige, T., & Skoog, F. (1962). A Revised Medium for Rapid Growth and Bio Assays with Tobacco Tissue Cultures. *Physiologia Plantarum*, 15(3), 473-497. DOI: 10.1111/j.1399-3054.1962.tb08052.x
- Quintero-Jiménez, A., Espinosa-Huerta, E., Acosta-Gallegos, J. A., Guzmán-Maldonado, H. S., & Mora-Avilés, M. A. (2010). Enhanced shoot organogenesis and regeneration in the common bean (*Phaseolus vulgaris* L.). *Plant Cell, Tissue and Organ Culture (PCTOC)*, 102(3), 381-386. DOI: 10.1007/s11240-010-9744-2
- Rech, E. L., Vianna, G. R., & Aragão, F. J. (2008). High-efficiency transformation by biolistics of soybean, common bean and cotton transgenic plants. *Nature Protocols*, 3(3), 410.
- Russell, D. R., Wallace, K. M., Bathe, J. H., Martinell, B. J., & McCabe, D. E. (1993). Stable transformation of *Phaseolus vulgaris* via electric-discharge mediated particle acceleration. *Plant Cell Reports*, 12(3), 165-169. DOI: 10.1007/BF00239099
- Santamaría, J. M., Hernández-Portilla, D., Chi-Manzanero, B., Espadas, F., Castaño, E., Iturriaga, G., & Rodríguez-Zapata, L. C. (2009). Incorporation of two trehalose biosynthetic genes in banana increases trehalose levels and protects the photosynthetic apparatus from salt-stress damage. *The Journal of Horticultural Science and Biotechnology*, 84(6), 665-671. DOI: 10.1080/14620316.2009.11512583
- Solis-Ramos, L. Y., González-Estrada, T., Andrade-Torres, A., Godoy-Hernández, G., & Castaño de la Serna, E. (2010). Endogenous GUS-like activity in *Capsicum chinense* Jacq. *Electronic Journal of Biotechnology*, 13(4), 2-3.
- Solis-Ramos, L. Y., González-Estrada, T., Nahuath-Dzib, S., Zapata-Rodríguez, L. C., & Castaño, E. (2009). Overexpression of *WUSCHEL* in *C. chinense* causes ectopic morphogenesis. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 96(3), 279-287. DOI: 10.1007/s11240-008-9485-7
- Solis-Ramos, L. Y., Valdez, M., Alvarado-Barrantes, R., Mora-Umaña, F., Hernández, E., Barboza, N., &

- Ramírez-Fonseca, P. (2015). Effect of Gamma Irradiation and Selection with Fungus Filtrate (*Rhizoctonia solani* Kuhn) on the in vitro Culture of Common Bean (*Phaseolus vulgaris*). *American Journal of Plant Science*, 6, 2672-2685. DOI: 10.4236/ajps.2015.616269
- Vianna, G. R., Albino, M. M. C., Dias, B. B. A., de Mesquita Silva, L., Rech, E. L., & Aragão, F. J. L. (2004). Fragment DNA as vector for genetic transformation of bean (*Phaseolus vulgaris* L.). *Scientia Horticulturae*, 99(3), 371-378. DOI: 10.1016/S0304-4238(03)00107-9
- Yamaguchi-Shinozaki, K., & Shinozaki, K. (1994). A novel cis-acting element in an *Arabidopsis* gene is involved in responsiveness to drought, low-temperature, or high-salt stress. *The Plant Cell*, 6(2), 251-264.
- Zambre, M., Goossens, A., Cardona, C., Montagu, M. V., Terryn, N., & Angenon, G. (2005). A reproducible genetic transformation system for cultivated *Phaseolus acutifolius* (tepary bean) and its use to assess the role of arcelins in resistance to the Mexican bean weevil. *Theoretical and Applied Genetics*, 110(5), 914-924. DOI: 10.1007/s00122-004-1910-7
- Zambre, M., Terryn, N., Clercq, J. D., Buck, S. D., Dillen, W., Montagu, M. V., Angenon, G. (2003). Light strongly promotes gene transfer from *Agrobacterium tumefaciens* to plant cells. *Planta*, 216(4), 580-586. DOI: 10.1007/s00425-002-0914-2
- Zhang, Z., Coyne, D. P., & Mitra, A. (1997). Factors Affecting *Agrobacterium*-mediated Transformation of Common Bean. *Journal of the American Society for Horticultural Science*, 122(3), 300-305.